

Drawdowns Preceding Rallies in the Brownian Motion Model

Olympia Hadjiliadis, Princeton University, Department of Electrical Engineering.
Jan Večeř, Columbia University, Department of Statistics.

This version: January 4, 2006

Abstract

We study drawdowns and rallies of Brownian motion. A rally is defined as the difference of the present value of the Brownian motion and its historical minimum, while the drawdown is defined as the difference of the historical maximum and its present value. This paper determines the probability that a drawdown of a units precedes a rally of b units. We apply this result to examine stock market crashes and rallies in the geometric Brownian motion model.

1 Introduction

In this paper we determine the probability that a drawdown precedes a rally in the Brownian motion model. The probabilities are computed by means of the distribution function of the random variables $y_{T_1(a)}^+$ and $y_{T_2(b)}^-$, where $y_{T_1(a)}^+$ represents the value of the upward rally when the drawdown process reaches the level a for the first time, and $y_{T_2(b)}^-$ represents the value of the drawdown when the upward rally process reaches the level b for the first time. Using the results of Taylor [11] and Lehoczky [4] concerning the distribution of a stopped drifted Brownian motion at the first time of the downfall of level a , we are able to show that the probability density functions of $y_{T_1(a)}^+$ and $y_{T_2(b)}^-$ are exponential, but with a discrete mass at 0. The paper concludes with a solution to the problem of computing the probability that a drop of $(100 \cdot \alpha)\%$ from the running maximum of a stock price occurs before a rally of $(100 \cdot \beta)\%$ from its running minimum, given that the stock price follows a geometric Brownian motion.

Risk management of drawdowns and portfolio optimization with drawdown constraints is becoming increasingly important among the practitioners. Chekhlov et. al. [2] studied drawdown measures in portfolio optimization. Magdon-Ismail et. al. [5] determined the distribution of the maximum drawdown of Brownian motion.

Our results are connected to a recent paper by Meilijson [6], where the results of Taylor [11] and Lehoczky [4] are used to derive the expected time to a given drawdown of Brownian motion, as well as the stationary distribution of the drawdown process. An alternative derivation of the above expected value based on the expected delay of the CUSUM stopping time, appears in Hadjiliadis & Moustakides [3]. The CUSUM stopping time was first proposed by Page [8] and was used subsequently as a means of detecting a regime change in the Brownian motion model (see Shiryaev [9], Beibel [1], and Moustakides [7]).

In the next section we first outline the main results of the paper. We then proceed to give an example of an application of these results in the examination of stock market crashes and rallies. We finally give some concluding remarks.

2 Results

Consider the process X_t with the following dynamics:

$$X_t = W_t + \gamma t,$$

where $\gamma \in \mathbb{R}$ and W_t is a standard Brownian motion process.

The quantity

$$X_t - \inf_{s \in [0, t]} X_s$$

measures the size of the rally comparing the present value of the process to its historical minimum, while the quantity

$$\sup_{s \in [0, t]} X_s - X_t$$

measures the size of the drawdown comparing the present value of the process to its historical maximum.

The aim of this section is to determine the probability that a drawdown of size a precedes a rally of size b . We introduce the stopping times:

$$T_1(a) = \inf\{t \geq 0 : \sup_{s \in [0, t]} X_s - X_t = a, \quad a \in \mathbb{R}_+\},$$

and

$$T_2(b) = \inf\{t \geq 0 : X_t - \inf_{s \in [0, t]} X_s = b, \quad b \in \mathbb{R}_+\}.$$

Consider the stopping time $T(a, b) = T_1(a) \wedge T_2(b)$. The stopping times $T_1(a)$ and $T_2(b)$ indicate respectively, the first time that the drawdown process reaches the critical level a , $T_1(a)$, and the first time the upward rally process reaches the critical level b , $T_2(b)$. In this section, we compute the probabilities of the events $\{T(a, b) = T_1(a)\}$, which represents the event that the drawdown of size a occurs before the rally of size b , and $\{T(a, b) = T_2(b)\}$, which represents the event that the rally of size b occurs before the drawdown of size a .

In order to simplify notation we introduce the following processes:

$$\begin{aligned} m_t^+ &:= \inf_{s \in [0, t]} X_s, \\ m_t^- &:= \inf_{s \in [0, t]} (-X_s) = - \sup_{s \in [0, t]} X_s, \\ y_t^+ &:= X_t - m_t^+, \\ y_t^- &:= -X_t - m_t^-. \end{aligned}$$

Using the above notation, the stopping times $T_1(a)$ and $T_2(b)$ become

$$\begin{aligned} T_1(a) &= \inf\{t \geq 0 : y_t^- = a, \quad a \in \mathbb{R}_+\} \\ T_2(b) &= \inf\{t \geq 0 : y_t^+ = b, \quad b \in \mathbb{R}_+\} \end{aligned}$$

Theorem 2.1 *Let $X_t = W_t + \gamma t$ be a standard Brownian motion with drift $\gamma \in \mathbb{R}$; T , T_1 and T_2 be the stopping times defined above. We distinguish the following two cases:*

1. $b \geq a > 0$

The probability of the drawdown preceding the rally, or the rally preceding the drawdown, are given respectively by

$$(2.1) \quad P\left(T(a, b) = T_1(a)\right) = m_A + (1 - m_A) \cdot \left[1 - \exp\left(-\frac{2\gamma}{e^{2\gamma a} - 1} \cdot (b - a)\right)\right],$$

$$(2.2) \quad P\left(T(a, b) = T_2(b)\right) = (1 - m_A) \cdot \exp\left(-\frac{2\gamma}{e^{2\gamma a} - 1} \cdot (b - a)\right),$$

where

$$(2.3) \quad m_A = \frac{e^{-2\gamma a} + 2\gamma a - 1}{e^{2\gamma a} + e^{-2\gamma a} - 2}.$$

2. $a \geq b > 0$

The probability of the drawdown preceding the rally, or the rally preceding the drawdown, are given respectively by

$$(2.4) \quad P\left(T(a, b) = T_1(a)\right) = (1 - m_B) \cdot \exp\left(-\frac{2\gamma}{1 - e^{-2\gamma b}} \cdot (a - b)\right),$$

$$(2.5) \quad P\left(T(a, b) = T_2(b)\right) = m_B + (1 - m_B) \cdot \left[1 - \exp\left(-\frac{2\gamma}{1 - e^{-2\gamma b}} \cdot (a - b)\right)\right],$$

where

$$(2.6) \quad m_B = \frac{e^{2\gamma b} - 2\gamma b - 1}{e^{2\gamma b} + e^{-2\gamma b} - 2}.$$

The proof of the theorem uses the following proposition:

Proposition 2.2 *The probability distribution functions of the random variables $y_{T_1(a)}^+$ and $y_{T_2(b)}^-$ are given by:*

1.

$$(2.7) \quad P(y_{T_1(a)}^+ = 0) = m_A,$$

$$(2.8) \quad P(y_{T_1(a)}^+ \in dr) = (1 - m_A) \cdot \left[\frac{2\gamma}{e^{2\gamma a} - 1} \cdot \exp\left(-\frac{2\gamma}{e^{2\gamma a} - 1} \cdot r\right)\right] dr, \quad r > 0,$$

where m_A is given by equation (2.3).

2.

$$(2.9) \quad P(y_{T_2(b)}^- = 0) = m_B,$$

$$(2.10) \quad P(y_{T_2(b)}^- \in dr) = (1 - m_B) \cdot \left[\frac{2\gamma}{1 - e^{-2\gamma b}} \cdot \exp\left(-\frac{2\gamma}{1 - e^{-2\gamma b}} \cdot r\right)\right] dr, \quad r > 0,$$

where m_B is given by equation (2.6).

In order to prove Proposition 2.2 and Theorem 2.1, we will need the following two lemmas.

Lemma 2.3 For $a, b \in \mathbb{R}_+$, we have:

$$(2.11) \quad E [T_1(a)] = \frac{e^{2\gamma a} - 2\gamma a - 1}{2\gamma^2},$$

$$(2.12) \quad E [T_2(b)] = \frac{e^{-2\gamma b} + 2\gamma b - 1}{2\gamma^2}.$$

PROOF. Let $g(x) = e^{-2\gamma x} + 2\gamma x - 1$. By applying Itô's rule to the process $g(y_t^+)$ we get

$$(2.13) \quad dg(y_t^+) = g'(y_t^+)dW_t + \gamma g'(y_t^+)dt - g'(y_t^+)dm_t^+ + \frac{1}{2}g''(y_t^+)dt.$$

We notice that the third term in the right hand side of the above equality disappears because $dm_t^+ \neq 0$ only when $y_t^+ = 0$ and $g'(0) = 0$. We also notice that the function g satisfies the second order differential equation

$$(2.14) \quad \gamma g'(x) + \frac{1}{2}g''(x) = 2\gamma^2.$$

Solving equation (2.13), we get:

$$(2.15) \quad g(y_t^+) - g(0) = \int_0^t g'(y_s^+)dW_s + \int_0^t \left(\gamma g'(y_s^+) + \frac{1}{2}g''(y_s^+) \right) ds$$

Let

$$S_n = \inf\{t \geq 0; 2\gamma^2 t \geq n\}.$$

Let $T_2^n(b) = T_2(b) \wedge S_n$. Obviously, $T_2^n(b)$ is *a.s.* finite. On the event $\{T_2^n(b) \geq t\}$, we have $\{y_t^+ \leq b\}$. Consequently,

$$(2.16) \quad E \left[\int_0^{T_2^n(b)} 2\gamma^2 g'(y_s^+) ds \right] \leq (g'(b))^2 n < \infty.$$

Evaluating (2.15) at $T_2^n(b)$, and taking expectations, while using equations (2.14) and (2.16) we get

$$E \left[g(y_{T_2^n(b)}^+) \right] = 2\gamma^2 E [T_2^n(b)].$$

But

$$(2.17) \quad g(b) \geq E \left[g(y_{T_2^n(b)}^+) \right].$$

Hence we have that

$$g(b) \geq 2\gamma^2 E [T_2^n(b)] \geq 2\gamma^2 E \left[1_{\{T_2^n(b) = \infty\}} \right].$$

Letting $n \rightarrow \infty$ and using the bounded convergence theorem, we deduce that $T_2(b)$ is finite *a.s.* as well.

Evaluating both sides of equation (2.15) at $T_2(b)$, and taking expectations of both sides, while using equation (2.14), and the fact that the quadratic variation of the stochastic integral $\int_0^{T_2(b)} g'(y_t^+)dW_t$ is finite (this follows from equation (2.16) and the *a.s.* finiteness of $T_2(b)$), we get

$$(2.18) \quad g(b) = 2\gamma^2 E [T_2(b)].$$

Consequently,

$$(2.19) \quad E[T_2(b)] = \frac{g(b)}{2\gamma^2}.$$

Similarly, we can show that

$$(2.20) \quad E[T_1(a)] = \frac{g(-a)}{2\gamma^2}.$$

This concludes the proof of the lemma. \diamond

Lemma 2.4 *We have*

$$y_t^+ + y_t^- = \max\left\{\sup_{s \leq t} y_s^+, \sup_{s \leq t} y_s^-\right\}.$$

PROOF. Observe that

$$(2.21) \quad y_t^+ + y_t^- = -m_t^+ - m_t^-.$$

We notice that the process $y_t^+ + y_t^-$ can only increase when either $X_t = m_t^+$ or $-X_t = m_t^-$, both of which cannot happen at the same time, since that would imply that $y_t^+ + y_t^-$ is 0. Therefore, $y_t^+ + y_t^-$ is a constant as a function of time unless either $y_t^+ = 0$ or $y_t^- = 0$, at which instant t , we simultaneously have:

1. $\max\{y_t^+, y_t^-\} = \max\left\{\sup_{s \leq t} y_s^+, \sup_{s \leq t} y_s^-\right\}$,
2. $\sup_{s \leq t} (y_s^+ + y_s^-) = y_t^+ + y_t^-$.

This completes the proof of the lemma. \diamond

As a consequence of this lemma we have

$$(2.22) \quad y_{T_1(a)}^+ = \left(\max_{t \leq T_1(a)} y_t^+ - a\right) \vee 0,$$

$$(2.23) \quad y_{T_2(b)}^- = \left(\max_{t \leq T_2(b)} y_t^- - b\right) \vee 0.$$

Finally, in order to proceed to the proof of Proposition 2.2 and Theorem 2.1, we will use the results of Taylor in [11] and Lehoczky in [4]. Taylor computes the bivariate Laplace transform of $X_{T_1(a)}$ and $T_1(a)$, where T_1 is defined as above. Lehoczky pointed out that the random variable $X_{T_1(a)} + a = \sup_{t \leq T_1(a)} X_t$ has the exponential distribution:

$$(2.24) \quad X_{T_1(a)} + a \sim \text{Exp}\left(\frac{2\gamma}{e^{2\gamma a} - 1}\right).$$

Note that the exponential parameter becomes equal to $\frac{1}{a}$ in the case when $\gamma = 0$. Now we can proceed to the proof of Proposition 2.2 and then to the proof of Theorem 2.1.

PROOF OF PROPOSITION 2.2. We will only compute the probability density function of the random variable $y_{T_1(a)}^+$ since the computation of the probability density function of the random variable $y_{T_2(b)}^-$ is done in a similar way. From equation (2.22), it follows that

$$(2.25) \quad P\left(y_{T_1(a)}^+ = 0\right) = P\left(\max_{t \leq T_1(a)} y_t^+ < a\right),$$

while

$$(2.26) \quad P\left(y_{T_1(a)}^+ \in dr\right) = P\left(\max_{t \leq T_1(a)} y_t^+ \geq a\right) \cdot P\left(y_{T_1(a)}^+ \in dr \mid \max_{t \leq T_1(a)} y_t^+ \geq a\right) \\ = P\left(y_{T_1(a)}^+ > 0\right) \cdot P\left(y_{T_1(a)}^+ \in dr \mid y_{T_1(a)}^+ > 0\right), \quad r > 0.$$

In the next discussion we first demonstrate

$$(2.27) \quad \mathcal{L}\left(y_{T_1(a)}^+ \mid y_{T_1(a)}^+ > 0\right) = \mathcal{L}\left(X_{T_1(a)} + a\right).$$

To see this, let

$$(2.28) \quad R_1 = \sup\{t \leq T_1(a); y_t^+ = 0\}.$$

Fix $r > 0$. Then

$$P\left(y_{T_1(a)}^+ \in dr \mid y_{T_1(a)}^+ > 0\right) = \frac{P\left(X_{T_1(a)} - \inf_{s \leq T_1(a)} X_s \in dr\right)}{P\left(\max_{t \leq T_1(a)} y_t^+ \geq a\right)} \\ = \frac{P\left(X_{T_1(a)} - X_{R_1} + X_{R_1} - \inf_{s \leq T_1(a)} X_s \in dr \mid R_1 < T_1(a)\right) \cdot P\left(R_1 < T_1(a)\right)}{P\left(\max_{t \leq T_1(a)} (X_t - X_{R_1} + X_{R_1} - \inf_{s \leq t} X_s) \geq a \mid R_1 < T_1(a)\right) \cdot P\left(R_1 < T_1(a)\right)} \\ = \frac{P\left(X_{T_1(a)} - X_{R_1} + X_{R_1} - \inf_{s \leq R_1} X_s \in dr \mid R_1 < T_1(a)\right)}{P\left(\max_{t \leq T_1(a)} (X_t - X_{R_1} + X_{R_1} - \inf_{s \leq R_1} X_s) \geq a \mid R_1 < T_1(a)\right)} = \frac{P\left(X_{T_1(a)} - X_{R_1} \in dr \mid R_1 < T_1(a)\right)}{P\left(\max_{t \leq T_1(a)} X_t - X_{R_1} \geq a \mid R_1 < T_1(a)\right)} \\ = \frac{P\left(X_{T_1(a)} - X_{R_1} \in dr \mid R_1 < T_1(a)\right)}{P\left(\max_{R_1 \leq t \leq T_1(a)} X_t - X_{R_1} \geq a \mid R_1 < T_1(a)\right)} = \frac{P(X_{T_1(a)} \in dr)}{P(\max_{t \leq T_1(a)} X_t \geq a)} \\ = \frac{\lambda e^{-\lambda r} e^{-\lambda a} dr}{e^{-\lambda a}} = \lambda e^{-\lambda r} dr = P(X_{T_1(a)} + a \in dr),$$

where $\lambda = \frac{2\gamma}{1 - e^{-2\gamma a}}$. Therefore, we get

$$(2.29) \quad P\left(y_{T_1(a)}^+ \in dr \mid y_{T_1(a)}^+ > 0\right) \sim \text{Exp}\left(\frac{2\gamma}{e^{2\gamma a} - 1}\right), \quad r > 0.$$

From equation (2.22), it follows that

$$(2.30) \quad P(y_{T_1(a)}^+ = 0) = P\left(T_1(a) < T_2(a)\right).$$

To compute $P\left(T_1(a) < T_2(a)\right)$, we first notice that

$$(2.31) \quad T_1(a) = T(a, b) + \left(T_1(a) - T(a, b)\right) \mathbf{1}_{\{T(a, b) = T_2(b)\}},$$

$$(2.32) \quad T_2(b) = T(a, b) + \left(T_2(b) - T(a, b)\right) \mathbf{1}_{\{T(a, b) = T_1(a)\}}.$$

Taking expectations we get

$$(2.33) \quad E[T_1(a)] = E[T(a, b)] + E\left[\left(T_1(a) - T(a, b)\right) \mathbf{1}_{\{T(a, b) = T_2(b)\}}\right],$$

$$(2.34) \quad E[T_2(b)] = E[T(a, b)] + E\left[\left(T_2(b) - T(a, b)\right) \mathbf{1}_{\{T(a, b) = T_1(a)\}}\right].$$

With $a = b$ and equation (2.22), it follows that

$$(2.35) \quad E [T_1(a)] = E [T(a, a)] + E [T_1(a)] \cdot P(T_2(a) < T_1(a)),$$

$$(2.36) \quad E [T_2(a)] = E [T(a, a)] + E [T_2(a)] \cdot P(T_1(a) < T_2(a)).$$

Using

$$P(T_1(a) < T_2(a)) + P(T_2(a) < T_1(a)) = 1$$

and equations (2.35) and (2.36), we conclude that

$$(2.37) \quad P(T_1(a) < T_2(a)) = \frac{E [T_2(a)]}{E [T_2(a)] + E [T_1(a)]}.$$

The result now follows by substituting (2.29), (2.30), (2.37) into equations (2.26), and (2.25), while using Lemma 2.3. This completes the proof of Proposition 2.2. \diamond

PROOF OF THEOREM 2.1. We will prove the theorem in the case that $b \geq a$ since the proof is similar in the case $a \geq b$. Suppose that $b \geq a$.

From Lemma 2.4 and equation (2.22), it follows that on the event $\{T_1(a) < T_2(b)\}$ we have

$$(2.38) \quad y_{T_1(a)}^+ = \begin{cases} 0 & \text{if } \max_{s \leq T_1(a)} y_s^+ < a, \\ \max_{s \leq T_1(a)} y_s^+ - a & \text{if } a \leq \max_{s \leq T_1(a)} y_s^+ < b. \end{cases}$$

Therefore,

$$(2.39) \quad P(T_1(a) < T_2(b)) = P(y_{T_1(a)}^+ = 0) + \int_{0^+}^{b-a} P(y_{T_1(a)}^+ \in dr),$$

and the result is obtained from Proposition 2.2. This completes the proof of Theorem 2.1. \diamond

Corollary 2.5 *Let $X_t = W_t$ be a standard Brownian motion and let T , T_1 and T_2 be stopping times defined as above. We distinguish the following two cases:*

1. $b \geq a > 0$

The probability of the drawdown preceding the rally, or the rally preceding the drawdown, are given respectively by

$$(2.40) \quad P(T(a, b) = T_1(a)) = \frac{1}{2} + \frac{1}{2} \cdot \left[1 - e^{-\frac{1}{a}(b-a)}\right],$$

$$(2.41) \quad P(T(a, b) = T_2(b)) = \frac{1}{2} \cdot e^{-\frac{1}{a}(b-a)}.$$

2. $a \geq b > 0$

The probability of the drawdown preceding the rally, or the rally preceding the drawdown, are given respectively by

$$(2.42) \quad P(T(a, b) = T_1(a)) = \frac{1}{2} \cdot e^{-\frac{1}{b}(a-b)},$$

$$(2.43) \quad P(T(a, b) = T_2(b)) = \frac{1}{2} + \frac{1}{2} \cdot \left[1 - e^{-\frac{1}{b}(a-b)}\right].$$

PROOF. It is a simple consequence of Theorem 2.1 by taking the limit as $\gamma \rightarrow 0$. \diamond

Corollary 2.6 *Let $X_t = W_t$ be a standard Brownian motion. The probability distribution function of the random variables $y_{T_1(a)}^+$ and $y_{T_2(b)}^-$ are given by*

1.

$$(2.44) \quad P(y_{T_1(a)}^+ = 0) = \frac{1}{2},$$

$$(2.45) \quad P(y_{T_1(a)}^+ \in dr) = \frac{1}{2} \cdot \left[\frac{1}{a} e^{-\frac{1}{a}r} \right] dr, \quad r > 0.$$

2.

$$(2.46) \quad P(y_{T_2(b)}^- = 0) = \frac{1}{2},$$

$$(2.47) \quad P(y_{T_2(b)}^- \in dr) = \frac{1}{2} \cdot \left[\frac{1}{b} e^{-\frac{1}{b}r} \right] dr, \quad r > 0.$$

PROOF. The above corollary is a consequence of Proposition 2.2 by letting $\gamma \rightarrow 0$. \diamond

Example: Stock Market Crashes and Rallies. Suppose that we have a stock S_t whose dynamics follow a geometric Brownian motion:

$$(2.48) \quad dS_t = \mu S_t dt + \sigma S_t dW_t.$$

What is the probability that this stock would drop by $(100 \cdot \alpha)\%$ before it incurs a rally of $(100 \cdot \beta)\%$? We can solve this problem by using Theorem 2.1. First observe that

$$(2.49) \quad S_t = S_0 \exp\left(\left(\mu - \frac{1}{2}\sigma^2\right)t + \sigma W_t\right) = S_0 \exp(\sigma X_t),$$

where $X_t = W_t + \left(\frac{\mu}{\sigma} - \frac{\sigma}{2}\right)t$. Let $U_1(\alpha)$ be the first time the stock drops by $(100 \cdot \alpha)\%$ from its running maximum and $U_2(\beta)$ be the first time the stock rallies by $(100 \cdot \beta)\%$ from its running minimum. Let

$$M_t = \sup_{0 \leq s \leq t} S_s,$$

and

$$N_t = \inf_{0 \leq s \leq t} S_s.$$

Notice that

$$(2.50) \quad \begin{aligned} U_1(\alpha) &= \inf\{t \geq 0 : S_t \leq (1 - \alpha)M_t\} \\ &= \inf\{t \geq 0 : \sup_{0 \leq s \leq t} X_s - X_t \geq -\frac{1}{\sigma} \log(1 - \alpha)\} = T_1\left(-\frac{1}{\sigma} \log(1 - \alpha)\right), \end{aligned}$$

and

$$(2.51) \quad \begin{aligned} U_2(\beta) &= \inf\{t \geq 0 : S_t \geq (1 + \beta)N_t\} \\ &= \inf\{t \geq 0 : X_t - \inf_{0 \leq s \leq t} X_s \geq \frac{1}{\sigma} \log(1 + \beta)\} = T_2\left(\frac{1}{\sigma} \log(1 + \beta)\right). \end{aligned}$$

Thus

$$(2.52) \quad P(U_1(\alpha) < U_2(\beta)) = P\left(T_1\left(-\frac{1}{\sigma} \log(1 - \alpha)\right) < T_2\left(\frac{1}{\sigma} \log(1 + \beta)\right)\right),$$

and we can apply Theorem 2.1 with the following parameters: $a = -\frac{1}{\sigma} \log(1 - \alpha)$, $b = \frac{1}{\sigma} \log(1 + \beta)$, $\gamma = \frac{\mu}{\sigma} - \frac{\sigma}{2}$. The resulting probability is given by:

$$(2.53) \quad P(U_1(\alpha) < U_2(\beta)) = \frac{(1+\beta)^{-\lceil 2\frac{\mu}{\sigma^2}-1 \rceil} + \left(2\frac{\mu}{\sigma^2}-1\right) \cdot \log(1+\beta) - 1}{(1+\beta)^{-\lceil 2\frac{\mu}{\sigma^2}-1 \rceil} + (1+\beta)^{\lceil 2\frac{\mu}{\sigma^2}-1 \rceil - 2}} \cdot [(1-\alpha)(1+\beta)]^{\left(\frac{2\frac{\mu}{\sigma^2}-1}{1-(1+\beta)^{\lceil 2\frac{\mu}{\sigma^2}-1 \rceil}}\right)},$$

when $\alpha \geq \frac{\beta}{1+\beta}$, and

$$(2.54) \quad P(U_1(\alpha) < U_2(\beta)) = 1 - \frac{(1-\alpha)^{-\lceil 2\frac{\mu}{\sigma^2}-1 \rceil} + \left(2\frac{\mu}{\sigma^2}-1\right) \cdot \log(1-\alpha) - 1}{(1-\alpha)^{-\lceil 2\frac{\mu}{\sigma^2}-1 \rceil} + (1-\alpha)^{\lceil 2\frac{\mu}{\sigma^2}-1 \rceil - 2}} \cdot [(1-\alpha)(1+\beta)]^{\left(\frac{2\frac{\mu}{\sigma^2}-1}{1-(1-\alpha)^{\lceil 2\frac{\mu}{\sigma^2}-1 \rceil}}\right)},$$

when $\alpha \leq \frac{\beta}{1+\beta}$. In the case when $\mu = \frac{1}{2}\sigma^2$, the above formulae simplify to

$$(2.55) \quad P(U_1(\alpha) < U_2(\beta)) = \frac{1}{2} [(1-\alpha)(1+\beta)]^{\frac{1}{\log(1+\beta)}},$$

when $\alpha \geq \frac{\beta}{1+\beta}$, and

$$(2.56) \quad P(U_1(\alpha) < U_2(\beta)) = 1 - \frac{1}{2} [(1-\alpha)(1+\beta)]^{\frac{1}{\log(1-\alpha)}},$$

when $\alpha \leq \frac{\beta}{1+\beta}$.

3 Conclusions

In this paper, we are able to compute the probability that a drawdown of size a occurs before an upward rally of size b in the drifted Brownian motion model and use this result to compute the probability that a drop of a given percentage α from the running maximum of a stock occurs before a rise of a given percentage β from the running minimum of a stock.

It is worth pointing out that using these results, one can compute the expected time of the minimum of the stopping times $T_1(a)$ and $T_2(b)$. This is of great interest for detecting a regime change in the case of two-sided alternatives. Stopping times such as the minimum of $T_1(a)$ and $T_2(b)$ resemble the 2-CUSUM stopping rule that has traditionally been used in the detection of regime changes in the presence of two-sided alternatives (see for example Hadjiliadis & Moustakides [3], Tartakovsky [10]).

References

- [1] M. Beibel. A note on Ritov's Bayes approach to the minimax property of the CUSUM procedure. *Annals of Statistics*, 24(4):1804–1812, 1996.
- [2] A. Chekhlov, S. Uryasev, and M. Zabarankin. Drawdown measure in portfolio optimization. *International Journal of Theoretical and Applied Finance*, 8(1):13–58, 2005.

- [3] O. Hadjiliadis and G. V. Moustakides. Optimal and asymptotically optimal CUSUM rules for change point detection in the Brownian motion model with multiple alternatives. *Theory of Probability and its applications (Teoriya Veroyatnostei i ee Primeneniya)*, 50(1):131–144, 2005.
- [4] J. P. Lehoczky. Formulas for stopped diffusion processes with stopping times based on the maximum. *Annals of Probability*, 5(4):601–607, 1977.
- [5] M. Magdon-Ismail, A. F. Atiya, A. Pratap, and Y. S. Abu-Mostafa. On the Maximum draw-down of a Brownian motion. *Journal of Applied Probability*, 41(1):147–161, 2004.
- [6] I. Meijilson. *The time to a given drawdown in Brownian motion*, volume XXXVII of *Seminaire de Probabilites*, pages 94–108. Springer Verlag, 2003.
- [7] G. V. Moustakides. Optimality of the CUSUM procedure in continuous time. *Annals of Statistics*, 32(1):302–315, 2004.
- [8] E. S. Page. Continuous inspection schemes. *Biometrika*, 41:100–115, 1954.
- [9] A. N. Shiryaev. Minimax optimality of the method of cumulative sums (CUSUM) in the case of continuous time. *Russian Mathematical Surveys*, 51:750–751, 1996.
- [10] A. G. Tartakovsky. Asymptotically minimax multi-alternative sequential rule for disorder detection. *Statistics and Control of Random Processes: Proceedings of the Steklov Institute of Mathematics*, 202(4):229–236, 1994. American Mathematical Society, Providence, Rhode Island.
- [11] H. M. Taylor. A stopped Brownian motion formula. *Annals of Probability*, 3(2):234–246, 1975.